

Laplacian spectral characterization of some double starlike trees

Pengli Lu¹ and Xiaogang Liu²

1. School of Computer and Communication, Lanzhou University of Technology, Lanzhou, 730050, Gansu, P.R. China.

2. Department of Mathematics and Statistics, The University of Melbourne, Parkville, VIC 3010, Australia

Email: lupengli88@163.com (P. Lu), lxg.666@163.com (X. Liu)

Abstract

A tree is called *double starlike* if it has exactly two vertices of degree greater than two. We denote by $H(p, n, q)$ one double starlike tree, which is obtained by attaching the centers of two stars $K_{1,p}$ and $K_{1,q}$ to the ends of a path P_n respectively. In this paper, graph $H(p, n, q)$ is proved to be determined by its Laplacian spectrum.

keywords: Laplacian spectrum; L -cospectral graphs; Line graph

AMS Subject Classification (2010): 05C50

1 Introduction

We start with some standard conceptions of graphs, as introduced in most textbooks on graph theory (e.g. [1, 3]). Let $G = (V(G), E(G))$ be an undirected and simple graph (loops and multiple edges are not allowed) with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$, where v_1, v_2, \dots, v_n are indexed in the non-increasing order of degrees. Let matrix $A(G)$ be the $(0,1)$ -adjacency matrix of G and $d_i = d_i(G) = d_G(v_i)$ the degree of the vertex v_i . The matrix $L(G) = D(G) - A(G)$ is called the *Laplacian matrix* of G , where $D(G)$ is the $n \times n$ diagonal matrix with $\{d_1, d_2, \dots, d_n\}$ as diagonal entries. The eigenvalues of matrices $A(G)$ and $L(G)$ are called the *adjacency eigenvalues* and *Laplacian eigenvalues* of G , respectively. We denote by $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ and $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n (= 0)$ the adjacency eigenvalues and the Laplacian eigenvalues of G , respectively. The multiset of eigenvalues of $A(G)$ (resp. $L(G)$) is called the *adjacency* (resp. *Laplacian*) *spectrum* of G . Two graphs are said to be *A-cospectral* (resp. *L-cospectral*) if they have the same adjacency (resp. Laplacian) spectrum.

A graph is said to be determined by its Laplacian (resp. adjacency) spectrum if there is no other non-isomorphic graphs L -cospectral (resp. A -cospectral) with it. To characterize some graphs by their spectra is very difficult. For its background, we refer the readers to [3–5].




Fig. 1. Double starlike tree $H(p, n, q)$

Up until now, some trees with special structures have been proved to be determined by their spectra, for example, the path [4], the T -shape tree [20], and the starlike tree [17]. Moreover, one double starlike tree, denoted by $H_n(p, p)$ ($n \geq 2, p \geq 1$), was proved to be determined by its

Laplacian spectrum in [11]. In the Conclusion of [11], the authors posed the double starlike tree $H(p, n, q)$ ($n \geq 1, p \geq 1, q \geq 1$) (shown in Figure 1) and indicated the difficulties to prove that $H(p, n, q)$ is determined by its Laplacian spectrum. Indeed, if we let $p = q \geq 2$, then $H(p, n, p)$ is $H_n(p, p)$. Clearly, if $p = q = 1$, then $H(1, n, 1)$ is a path of order $n + 2$. If $n = 1$, then $H(p, 1, q)$ is a star $K_{1, p+q}$ of order $p + q + 1$. It is known that paths, stars and $H_n(p, p)$ are all determined by their Laplacian spectra [4, 11, 17]. Therefore, we will only need to consider $H(p, n, q)$ for $n \geq 2, p \neq q \geq 2$. Without loss of generality, we let $p > q \geq 2$. In [13–15, 19], the authors investigated double starlike trees $H(p, n, q)$ with $p > q = 2$ or $p - q = 1, 2, 7$ and proved that they were determined by their Laplacian spectra respectively. Unfortunately, they did not present the most powerful method to solve the problem of Laplacian spectral characterization of $H(p, n, q)$, although they solved it partially.

In this paper, we will solve this problem completely. We pose a new method for $H(p, n, q)$ with $n \geq 2, p > q \geq 2$. Then together with the result in [11], we can conclude that every double starlike tree $H(p, n, q)$ is determined by its Laplacian spectrum.

2 Preliminaries

In order to show our main result, some previously established results are summarized in this section.

Lemma 2.1. [6] *Let T be a tree with n vertices and $\mathcal{L}(T)$ be its line graph. Then for $i = 1, 2, \dots, n - 1$, $\mu_i(T) = \lambda_i(\mathcal{L}(T)) + 2$.*

Lemma 2.2. [12] *Let u be a vertex of G , and $G - u$ be the subgraph obtained from G by deleting u together with its incident edges. Then*

$$\mu_i(G) \geq \mu_i(G - u) \geq \mu_{i+1}(G) - 1, \quad i = 1, 2, \dots, n - 1.$$

Lemma 2.3. [4] *Suppose that N is a symmetric $n \times n$ matrix with eigenvalues $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n$. Then the eigenvalues $\alpha'_1 \geq \alpha'_2 \geq \dots \geq \alpha'_m$ of a principal submatrix of N of size m satisfy $\alpha_i \geq \alpha'_i \geq \alpha_{n-m+i}$ for $i = 1, 2, \dots, m$.*

Lemma 2.4. [2] *In a simple graph, the number of closed walks of length 4 equals twice the number of edges plus four times of the number of paths on three vertices plus eight times of the number of 4-cycles.*

Lemma 2.5. [4, 16] *Let G be a graph. For the adjacency matrix and the Laplacian matrix, the following can be deduced from the spectrum.*

(1) *The number of vertices.*

(2) *The number of edges.*

For the adjacency matrix, the following follows from the spectrum.

(3) *The number of closed walks of any length.*

For the Laplacian matrix, the following follows from the spectrum.

(4) *The number of components.*

(5) *The number of spanning trees.*

(6) *The sum of the squares of degrees of vertices.*

Lemma 2.6. [8,9] Let G be a graph with $V(G) \neq \emptyset$ and $E(G) \neq \emptyset$. Then

$$d_1 + 1 \leq \mu_1 \leq \max \left\{ \frac{d_i(d_i + m_i) + d_j(d_j + m_j)}{d_i + d_j}, \quad v_i v_j \in E(G) \right\},$$

where m_i denotes the average of the degrees of the vertices adjacent to vertex v_i in G .

Lemma 2.7. [10] Let G be a connected graph with $n \geq 3$ vertices. Then $\mu_2 \geq d_2$.

Lemma 2.8. [7] Let G be a connected graph with $n \geq 4$ vertices. Then $\mu_3 \geq d_3 - 1$.

Lemma 2.9. [3] If e is an edge of the graph G and $G' = G - e$. Then

$$\mu_1(G) \geq \mu_1(G') \geq \mu_2(G) \geq \mu_2(G') \geq \cdots \geq \mu_{n-1}(G) \geq \mu_{n-1}(G') \geq \mu_n(G) = \mu_n(G') = 0.$$

3 Main results

First, some properties on the Laplacian eigenvalues of the double starlike tree $H(p, n, q)$ are stated as follows:

Lemma 3.1. If $n \geq 4$ and $p > q \geq 2$, then the Laplacian eigenvalues of $H(p, n, q)$ satisfy

- (1) $p + 2 \leq \mu_1(H(p, n, q)) \leq p + 2 + \frac{1}{p + 2}$.
- (2) $q + 2 \leq \mu_2(H(p, n, q)) \leq q + 3 + \frac{1}{q + 2}$.
- (3) $\mu_3(H(p, n, q)) < 4$.

Proof. (1) It follows from Lemma 2.6 by a simple calculation.

(2) Let u and v be the vertices of degree $p + 1$ and $q + 1$ in $H(p, n, q)$, respectively. Then by Lemma 2.2, we have

$$\mu_1(H(p, n, q)) \geq \mu_1(H(p, n, q) - u) \geq \mu_2(H(p, n, q)) - 1.$$

Lemma 2.6 implies that

$$\mu_1(H(p, n, q) - u) \leq q + 2 + \frac{1}{q + 2}.$$

Then

$$\mu_2(H(p, n, q)) \leq \mu_1(H(p, n, q) - u) + 1 \leq q + 3 + \frac{1}{q + 2}.$$

Next, let G_1 be a subgraph of $H(p, n, q)$ obtained by deleting an edge whose end is neither u nor v . Clearly, G_1 has two connected components. Moreover, by Lemma 2.6, the largest Laplacian eigenvalue of each component is at least $q + 2$, i.e., $\mu_2(G_1) \geq q + 2$. Hence by Lemma 2.9, $\mu_2(H(p, n, q)) \geq \mu_2(G_1) \geq q + 2$.

(3) Let M_{uv} be the $(p + n + q - 2) \times (p + n + q - 2)$ principal submatrix of $L(H(p, n, q))$ formed by deleting the rows and columns corresponding to u and v . Clearly, the largest eigenvalue of M_{uv} is less than 4. By Lemma 2.3, $\mu_3(H(p, n, q)) < 4$. \square

Lemma 3.2. *Every graph L -cospectral to $H(p, n, q)$ with $n \geq 4$ and $p > q \geq 2$ is a tree with the degree sequence $(p+1, q+1, \overbrace{2, \dots, 2}^{n-2}, \overbrace{1, \dots, 1}^{p+q})$.*

Proof. Suppose that G' is L -cospectral to $H(p, n, q)$ with $n \geq 4$ and $p > q \geq 2$. By (1), (2), (4) and (5) of Lemma 2.5, G' is a tree with $n + p + q$ vertices and $n + p + q - 1$ edges. Let $d_1 \geq d_2 \geq \dots \geq d_{n+p+q}$ be the non-increasing degree sequence of graph G' . Now we determine the degree sequence of G' . By (1) of Lemma 3.1, we have $d_1 + 1 \leq \mu_1(G') \leq p + 2 + \frac{1}{p+2}$. Hence the largest degree $\Delta' = \Delta(G')$ of G' is at most $p+1$. Denote by n_i the number of vertices in G' with degree i , for $i = 1, 2, \dots, \Delta' = d_1 \leq p+1$. By (1), (2) and (6) of Lemma 2.5, we have

$$\sum_{i=1}^{\Delta'} n_i = n + p + q, \quad (3.1)$$

$$\sum_{i=1}^{\Delta'} i n_i = 2(n + p + q - 1), \quad (3.2)$$

$$\sum_{i=1}^{\Delta'} i^2 n_i = (p+1)^2 + (q+1)^2 + 4(n-2) + p + q. \quad (3.3)$$

By (3.1), (3.2) and (3.3), we have

$$\sum_{i=1}^{\Delta'} (i^2 - 3i + 2) n_i = p^2 + q^2 - p - q. \quad (3.4)$$

By Lemma 2.8 and (3) of Lemma 3.1, we have $d_3 \leq \mu_3(G') + 1 = \mu_3(H(p, n, q)) + 1 < 5$, which implies $d_3 \leq 4$. By Lemma 2.7 and (2) of Lemma 3.1, we have $d_2 \leq \mu_2(G') = \mu_2(H(p, n, q)) \leq q + 3 + \frac{1}{q+2}$, which implies $d_2 \leq q + 3$. Lemma 2.1 implies that the line graphs of $\mathcal{L}(G')$ and $\mathcal{L}(H(p, n, q))$ are cospectral with respect to adjacency matrix, and by (3) of Lemma 2.5, they have the same number of triangles (sixth times of the number of closed walks of length 3), *i.e.*,

$$\sum_{i=1}^{\Delta'} \binom{i}{3} n_i = \binom{p+1}{3} + \binom{q+1}{3}. \quad (3.5)$$

Note that $\Delta' (= d_1) \leq p+1$, $d_2 \leq q+3$. We consider the following two cases.

Case 1. $q = 2$ or $q = 3$.

Case 1.1. $\Delta' (= d_1) < p+1$, *i.e.*, $n_{p+1} = 0$. By (3.5) and (3.4),

$$\binom{p+1}{3} + \binom{q+1}{3} = \sum_{i=1}^{\Delta'} \binom{i}{3} n_i \leq \frac{p}{6} \sum_{i=1}^{\Delta'} (i-1)(i-2) n_i = \frac{p}{6} (p^2 + q^2 - p - q).$$

If $q = 2$, we can get $p^2 - 3p + 6 \leq 0$, which is a contradiction to $p \geq 3$. If $q = 3$, we can get $p^2 - 7p + 24 \leq 0$, which is a contradiction to $p \geq 4$. Therefore $n_{p+1} \geq 1$.

Case 1.2. $n_{p+1} \geq 2$, i.e., there are at least 2 vertices with the largest degree $\Delta' = p + 1$. Then by (3.5),

$$\binom{p+1}{3} + \binom{q+1}{3} = \sum_{i=1}^{\Delta'} \binom{i}{3} n_i \geq 2 \binom{p+1}{3} + \sum_{i=1}^p \binom{i}{3} n_i,$$

i.e.,

$$\binom{q+1}{3} - \binom{p+1}{3} \geq \sum_{i=1}^p \binom{i}{3} n_i.$$

This is a contradiction to $p > q \geq 2$. Hence $n_{p+1} = 1$. Further, if $q = 2$, by (3.5), we have $n_i = 0$ for $i = 4, \dots, p$ and $n_3 = 1$. Moreover, by (3.1) and (3.2), we have $n_2 = n - 2$ and $n_1 = p + 2$. So the claim holds. If $q = 3$, by (3.5), we have $n_i = 0$ for $i = 5, \dots, p$ and $n_4 \leq 1$. By (3.1), (3.2) and (3.3), it is easy to see that $n_4 = 1$. Moreover, by (3.1), (3.2) and (3.3), $n_3 = 0$, $n_2 = n - 2$ and $n_1 = p + 3$. So the claim holds.

Case 2. $q \geq 4$. Clearly, $d_1 \geq 4$ (Otherwise, if $d_1 = 3$, by (3.4), $n_3 = \frac{1}{2}(p^2 + q^2 - p - q)$, which is a contradiction to (3.5)). By (3.4) and (3.5), we have

$$\sum_{i=1}^{\Delta'} (i-1)(i-2)(i-3)n_i = p^3 + q^3 - 3p^2 - 3q^2 + 2p + 2q.$$

So,

$$6n_4 + \sum_{i=5}^{\Delta'} (i-1)(i-2)(i-3)n_i = p(p-1)(p-2) + q(q-1)(q-2). \quad (3.6)$$

Note that $d_3 \leq 4$, which implies that G' has at most two vertices of degree strictly greater than 4. Consider the following three cases.

Case 2.1. $d_1 = 4$, $d_2 \leq 4$. By (3.6),

$$6n_4 = p(p-1)(p-2) + q(q-1)(q-2).$$

By (3.1), (3.2) and (3.3), we have

$$n_3 = -\frac{1}{2}p(p-1)(p-3) - \frac{1}{2}q(q-1)(q-3) < 0,$$

a contradiction.

Case 2.2. $d_1 > 4$, $d_2 = 4$. By (3.6),

$$6n_4 = p(p-1)(p-2) + q(q-1)(q-2) - (d_1-1)(d_1-2)(d_1-3).$$

By (3.1), (3.2) and (3.3), we have

$$n_3 = \frac{1}{2}(d_1-1)(d_1-2)(d_1-4) - \frac{1}{2}p(p-1)(p-3) - \frac{1}{2}q(q-1)(q-3).$$

For $d_1 \leq p+1$ and $q \geq 4$, $n_3 < 0$, a contradiction.

Case 2.3. $d_1 > 4$, $d_2 > 4$. By (3.6),

$$n_4 = \binom{p}{3} + \binom{q}{3} - \binom{d_1-1}{3} - \binom{d_2-1}{3}. \quad (3.7)$$

By (3.1), (3.2), (3.3) and (3.7), we have

$$n_3 = \frac{1}{2}(d_1-1)(d_1-2)(d_1-4) - \frac{1}{2}p(p-1)(p-3) + \frac{1}{2}(d_2-1)(d_2-2)(d_2-4) - \frac{1}{2}q(q-1)(q-3). \quad (3.8)$$

Case 2.3.1. $\Delta' (= d_1) < p+1$, i.e., $n_{p+1} = 0$. Then consider the following cases.

Case 2.3.1.1. $\Delta' (= d_1) = p$, $d_2 = q+3$. By (3.7) and (3.8), we have

$$n_4 = \frac{1}{2}p^2 - \frac{3}{2}p + 1 - q^2,$$

and

$$n_3 = -\frac{3}{2}p^2 + \frac{11}{2}p - 5 + 3q^2 - 2q = -\frac{1}{2}(p-2)(3p-5) + 3q^2 - 2q.$$

Then $n_4 \geq 0$ implies that $p \geq \frac{3 + \sqrt{8q^2 + 1}}{2}$. So,

$$n_3 \leq -\frac{1}{2} \left(\frac{3 + \sqrt{8q^2 + 1}}{2} - 2 \right) \left(3 \times \frac{3 + \sqrt{8q^2 + 1}}{2} - 5 \right) + 3q^2 - 2q = -\frac{1}{2} + \frac{1}{2}\sqrt{8q^2 + 1} - 2q < 0,$$

a contradiction.

Case 2.3.1.2. $\Delta' (= d_1) = p-1$, $d_2 = q+3$. By (3.7) and (3.8), we have

$$n_4 = p^2 - 4p - q^2 + 4 = (p-2-q)(p-2+q),$$

and

$$n_3 = -3p^2 + 14p - 16 + 3q^2 - 2q = -(p-2)(3p-8) + 3q^2 - 2q.$$

Then $n_4 \geq 0$ implies that $p \geq q+2$. So,

$$n_3 \leq -q(3q-2) + 3q^2 - 2q = 0.$$

Therefore $p = q+2$, and $n_3 = n_4 = 0$. But now $d_1 = p-1 = q+1 < d_2 = q+3$, a contradiction.

Case 2.3.1.3. $\Delta' (= d_1) \leq p-2$, $d_2 = q+3$. Then $q+3 = d_2 \leq d_1 \leq p-2$ implies that $p \geq q+5$. By (3.8), we have

$$n_3 \leq -\frac{9}{2}p^2 + \frac{51}{2}p - 37 + 3q^2 - 2q \leq -\frac{9}{2}(q+5)^2 + \frac{51}{2}(q+5) - 37 + 3q^2 - 2q \leq -\frac{3}{2}q^2 - \frac{43}{2}q - 22 < 0.$$

It is a contradiction.

Case 2.3.1.4. $\Delta' (= d_1) \leq p$, $d_2 \leq q + 2$. By (3.8) and $p \geq q + 1$, we have

$$n_3 \leq -\frac{3}{2}p^2 + \frac{11}{2}p - 4 + \frac{3}{2}q^2 - \frac{5}{2}q = -\frac{1}{2}(p-1)(3p-8) + \frac{3}{2}q^2 - \frac{5}{2}q \leq -\frac{1}{2}q(3q-5) + \frac{3}{2}q^2 - \frac{5}{2}q = 0.$$

Therefore $n_3 = 0$ implies that $d_1 = p$, $d_2 = q + 2$ and $p = q + 1$. But now $d_1 = p = q + 1 < q + 2 = d_2$, a contradiction.

Case 2.3.2. $\Delta' (= d_1) = p + 1$, i.e., $n_{p+1} \geq 1$. If $n_{p+1} \geq 2$, i.e., there are at least 2 vertices with degree $\Delta' = p + 1$, then by (3.5),

$$\binom{p+1}{3} + \binom{q+1}{3} = \sum_{i=1}^{\Delta'} \binom{i}{3} n_i \geq 2 \binom{p+1}{3} + \sum_{i=1}^p \binom{i}{3} n_i,$$

i.e.,

$$\binom{q+1}{3} - \binom{p+1}{3} \geq \sum_{i=1}^p \binom{i}{3} n_i.$$

This is a contradiction to $p > q \geq 4$. Hence $n_{p+1} = 1$. For $q \geq 4$, by (3.7) and (3.8), it is easy to see that $d_2 = q + 1$. By (3.5), $n_i = 0$ for $i = 3, 4, \dots, q, q+2, \dots, p$. By (3.1) and (3.2), $n_1 = p + q$

and $n_2 = n - 2$. Therefore, the degree sequence of G' is $(p + 1, q + 1, \overbrace{2, \dots, 2}^{n-2}, \overbrace{1, \dots, 1}^{p+q})$. \square

Now we can present the main result of this paper.

Theorem 3.3. *Every double starlike tree $H(p, n, q)$ is determined by its Laplacian spectrum.*

Proof. If $n = 1$, $H(p, 1, q)$ is a star $K_{1, p+q}$, which is determined by its Laplacian spectrum [17]. If $n = 2$ or $n = 3$, $H(p, n, q)$ is determined by their Laplacian spectra [18]. If $p = q = 1$, $H(1, n, 1)$ is a path, which is determined by its Laplacian spectrum [4]. If $p > q = 1$, $H(p, n, 1)$ is a starlike tree, which is determined by its Laplacian spectrum [17]. If $p = q \geq 2$, $H(p, n, p)$ is determined by its Laplacian spectrum [11]. Hence we only need to consider $n \geq 4$, $p > q \geq 2$. Suppose that G' is L -cospectral to $H(p, n, q)$ with $n \geq 4$ and $p > q \geq 2$. By Lemma 3.2, G' is a

tree with the degree sequence of G' is $(p + 1, q + 1, \overbrace{2, \dots, 2}^{n-2}, \overbrace{1, \dots, 1}^{p+q})$. In the following, we show that G' is isomorphic to $H(p, n, q)$.

First, we assume that the vertex of degree $p + 1$ is adjacent to the vertex of degree $q + 1$ and let a and b be such that there are $q - a$ (resp. $p - b$) vertices of degree 1 adjacent to the vertex of degree $q + 1$ (resp. $p + 1$) in G' (see Fig. 2); we have $0 \leq a \leq q$ and $0 \leq b \leq p$. Then,

$$\sum_{i=1}^a l'_i + \sum_{j=1}^b l''_j + (p + q) + 2 = n + p + q.$$

That is,

$$\sum_{i=1}^a l'_i + \sum_{j=1}^b l''_j = n - 2.$$



$(p+1, q+1, \overbrace{p, \dots, p}^p, \overbrace{q, \dots, q}^q, \overbrace{2, \dots, 2}^{n-3})$. By (3) of Lemma 2.5, $\mathcal{L}(H(p, n, q))$ and $\mathcal{L}(G')$ have the same number of closed walks of length 4. Clearly, $\mathcal{L}(H(p, n, q))$ and $\mathcal{L}(G')$ have the same number of edges. Moreover, they have the same number of 4-cycles. Hence by Lemma 2.4, $\mathcal{L}(H(p, n, q))$ and $\mathcal{L}(G')$ have the same number of paths on three vertices. Therefore,

Hence,

$$(p-1)(q-1) + b(p-1) + a(q-1) = 0.$$

But for $p > q \geq 2$, $0 \leq a \leq q$ and $0 \leq b \leq p$, we have

$$(p-1)(q-1) + b(p-1) + a(q-1) \neq 0.$$

It is a contradiction.

Now assume that the vertex of degree $p + 1$ is not adjacent to the vertex of degree $q + 1$ and let a and b be such that there are $q - a$ (resp. $p - b$) vertices of degree 1 adjacent to the vertex of degree $q + 1$ (resp. $p + 1$) in G' (see Fig. 3); we have $0 \leq a \leq q$ and $0 \leq b \leq p$. Then,

$$l_1 + \sum_{i=1}^a l'_i + \sum_{j=1}^b l''_j + (p+q) = n + p + q.$$

That is,

$$l_1 + \sum_{i=1}^a l'_i + \sum_{j=1}^b l''_j = n. \quad (3.9)$$



For the line graph $\mathcal{L}(G')$, suppose that there exist n'_i vertices of degree i . Then $n'_1 = a + b$, $n'_{p+1} = b + 1$, $n'_p = p - b$, $n'_{q+1} = a + 1$, $n'_q = q - a$, $n'_2 = n + p + q - 1 - (n'_1 + n'_{p+1} + n'_p +$

$n'_{q+1} + n'_q = n - 3 - a - b$, and $n'_j = 0$, where $j \notin \{1, 2, p, q, p+1, q+1\}$. By (3) of Lemma 2.5, $\mathcal{L}(H(p, n, q))$ and $\mathcal{L}(G')$ have the same number of closed walks of length 4. Clearly, $\mathcal{L}(H(p, n, q))$ and $\mathcal{L}(G')$ have the same number of edges and the same number of 4-cycles. Hence by Lemma 2.4, $\mathcal{L}(H(p, n, q))$ and $\mathcal{L}(G')$ have the same number of paths on three vertices. Therefore,

$$\begin{aligned} & \binom{p+1}{2} + \binom{q+1}{2} + p \binom{p}{2} + q \binom{q}{2} + (n-3) \binom{2}{2} \\ &= (b+1) \binom{p+1}{2} + (p-b) \binom{p}{2} + (a+1) \binom{q+1}{2} + (q-a) \binom{q}{2} + (n-3-a-b) \binom{2}{2}. \end{aligned}$$

Hence,

$$\frac{1}{2}p^3 + \frac{1}{2}q^3 + \frac{1}{2}p + \frac{1}{2}q + n - 3 = \frac{1}{2}p^3 + \frac{1}{2}q^3 + \frac{1}{2}p + \frac{1}{2}q + n - 3 + a(q-1) + b(p-1).$$

So $a(q-1) + b(p-1) = 0$, i.e., $a = 0$ and $b = 0$. By (3.9), we have $l_1 = n$. Hence, all of the vertices of degree 1 are adjacent to the vertices of degree $p+1$ and the vertices of degree $q+1$ in the graph G' . Then G' is isomorphic to $H(p, n, q)$. \square

Corollary 3.4. *The complement of a double starlike tree $H(p, n, q)$ is determined by its Laplacian spectrum.*

Proof. It follows from [8] and Theorem 3.3. \square

Acknowledgements P. Lu is supported by the National Natural Science Foundation of China (61064011) and the China Postdoctoral Science Foundation (20100470088). X. Liu is supported by MIFRS and MIRS of the University of Melbourne.

References

- [1] D.M. Cvetković, M. Doob, H. Sachs, Spectra of Graphs: Theory and Applications, Academic press, New York, San Francisco, London, 1980.
- [2] D.M. Cvetković, P. Rowlinson, Spectra of unicyclic graphs, Graphs Combin. 3 (1987) 7-23.
- [3] D.M. Cvetković, P. Rowlinson, S. Simić, An introduction to the theory of graph spectra, Cambridge University Press, Cambridge, 2010.
- [4] E.R. van Dam, W.H. Haemers, Which graphs are determined by their spectrum? Linear Algebra Appl. 373 (2003) 241-272.
- [5] E.R. van Dam, W.H. Haemers, Developments on spectral characterizations of graphs, Discrete Math. 309 (2009) 576-586.
- [6] I. Gutman, V. Gineityte, M. Lepović, M. Petrović, The high-energy band in the photoelectron spectrum of alkaners and its dependence on molecular structure, J. Serb. Chem. 64 (1999) 673-680.
- [7] J.-M. Guo, On the third largest Laplacian eigenvalue of a graph, Linear and Multilinear Algebra 55(1) (2007) 93-102.
- [8] A.K. Kelmans, V.M. Chelnokov, A certain polynomial of a graph and graphs with an extremal numbers of trees, J. Combin. Theory Ser. B 16 (1974) 197-214.
- [9] J.-S. Li, X.-D. Zhang, On the Laplacian eigenvalues of a graph, Linear Algebra Appl. 285 (1998) 305-307.
- [10] J.-S. Li, Y.-L. Pan, A note on the second largest eigenvalue of the Laplacian matrix of a graph, Linear and Multilinear Algebra 48(20) (2000) 117-121.
- [11] X.-G. Liu, Y.-P. Zhang, P.-L. Lu, One special double starlike graph is determined by its Laplacian spectrum, Applied Mathematics Letters, 22 (2009) 435-438.
- [12] Z. Lotker, Note on deleting a vertex and weak interlacing of the Laplacian spectrum, Electronic J. Linear Algebra, 16 (2007) 68-72.

- [13] P.-L. Lu, Z.-T. Yuan, Double-starlike tree $H_n(p, 2)$ determined by its Laplacian spectrum (in Chinese), Journal of Lanzhou University of Technology 34(2) (2008) 147-150.
- [14] P.-L. Lu, Z.-T. Yuan, Double Starlike Graphs with Largest and Second Largest Degree Difference 2 Determined by Their Laplacian Spectrum, CSIE 3 (2009) 26-29.
- [15] P.-L. Lu, X.-D. Zhang, Y.-P. Zhang, Determination of double quasi-star tree from its Laplacian spectrum, Journal of Shanghai University (English Edition), 14(3) (2010) 163-166.
- [16] C.S. Oliveira, N.M.M. de Abreu, S. Jurkiewicz, The characteristic polynomial of the Laplacian of graphs in (a, b) -linear cases, Linear Algebra Appl. 365 (2002) 113-121.
- [17] G.R. Omid, K. Tajbakhsh, Starlike trees are determined by their Laplacian spectrum, Linear Algebra Appl. 422 (2007) 654-658.
- [18] X.-L. Shen, Y.-P. Hou, Some trees are determined by their Laplacian spectra (in Chinese), Journal of Natural Science of Hunan Normal University 29(1) (2006) 21-24.
- [19] X.-L. Shen, Y.-P. Hou, Double-starlike Tree $D_{n,p,p-7}$ is Determined by Its Spectrum (in Chinese), Journal of Hunan University of Arts and Science (Natural Science Edition) 19(4) (2007) 21-24.
- [20] W. Wang, C.-X. Xu, Note: The T -shape tree is determined by its Laplacian spectrum, Linear Algebra Appl. 419 (2006) 78-81.